

FOAM product	indication	dose regime	DOSE SIZE	Reference drug (not foam)	Dose size	Dosage form	Dose regimen
Cortifoam Hydrocortisone acetate 10%	Ulcerative proctitis	once or twice daily for 2-3 weeks	Each application delivers approximately 900 mg of foam containing 90 mg of hydrocortisone acetate	CORTENEMA (Hydrocortisone 0.17%)	The usual dose is one 60 mL enema (100 mg hydrocortisone) daily for 2 or 3 weeks	enema	The usual course of therapy is one CORTENEMA* nightly for 21 days
Proctofoam HC Hydrocortisone acetate 1% Pramoxine hydrochloride 1%	Proctitis	Insert 1 applicatorful into anus 2 or 3 times daily and after each bowel evacuation	One application contains 375 mg of mucoadhesive base containing 1% hydrocortisone acetate USP (3.75 mg/dose) and 1% pramoxine hydrochloride USP (3.75 mg/dose).	Proctofoam (Hydrocortisone Cream, USP) 1%	Topical application to the affected area (no specified dose)	cream	Apply to the affected area as a thin film from two to four times daily depending on the severity of the condition.
Clobetasol Propionate FOAM 0.05%	psoriasis	Clobetasol propionate foam, 0.05% should be applied to the affected area twice daily, once in the morning and once at night	Topical application	Clobetasol Propionate 0.05%	Topical application	cream or ointment	Apply a thin layer of clobetasol propionate cream or ointment to the affected skin areas twice daily and rub in gently and completely

Extina Foam (ketoconazole 2%)	seborrheic dermatitis	Twice daily for 4 weeks	Topical application	Ketoconazole Gel 2%	Topical application	Gel	XOLEGEL should be applied <u>once daily</u> to the affected area for 2 weeks.
				Ketoconazole cream 2%	Topical application	cream	Ketoconazole Cream 2% should be applied to the affected area twice daily for four weeks or until clinical clearing
Luxiq Foam (Bethamethasone valerate 0.12%)	scalp psoriasis	Apply twice daily, once in the morning and once at night	Topical application	Betamethasone Valerate cream or ointment (0.1%)	Topical application	cream or ointment	Dosage once or twice a day is often effective
Verdeso Foam (desonide 0.05%)	moderate atopic dermatitis	A thin layer of Verdeso Foam should be applied to the affected area(s) twice daily	Topical application	Desonate (Desonide Gel 0.05%)	Topical application	Gel	Apply a thin layer to the affected areas two times daily
SORILUX Foam (calcipotriene 0.005%)	plaque psoriasis	Apply a thin layer of SORILUX Foam twice daily to the affected areas	Topical application	Dovonex Cream (calcipotriene cream), 0.005%	Topical application	Cream	Once or twice daily